

台中榮總小兒科常用水劑藥品使用劑量表(I)

藥品名稱	成份含量	一般劑量
* Acetaminophen <i>Anti-phen syrup</i>	24 mg/ml, 60ml/bot	Fever: 10-15 mg/kg/dose PO Q4H-Q6H, maximum 5 doses/day
Ambroxol <i>Losolvan solution</i>	3 mg/ml, 60 ml/bot	< 2 yrs: 1.2-1.6 mg/kg/day (分2次服用) 2-5 yrs: 2.5 ml BID
* Amoxicillin <i>Supercillin susp</i>	25 mg/ml, 60 ml/bot	20-40 mg/kg/day, Q6H 分次服用
*Amoxicillin 400 mg + Potassium clavulanate 57mg <i>Augmentin syrup</i>	457 mg/5ml, 35 ml/bot	2-6 yrs: 2.5ml Q12H 7-12 yrs: 5ml Q12H Sinusitis: 80-90 mg/kg/day 分2次服用
*Azithromycin <i>Zithromax susp</i>	200 mg/5 ml, 15 ml/bot	<15 kg: 10 mg/kg for 3 days 15-25 kg: 200mg (5ml)/QD for 3day
* Bromhexine <i>Bisolvon solution</i>	2 mg/ml, 50 ml/bot	<i>Infant</i> : 0.5ml TID 5 yrs : 1ml TID 5-10 yrs: 2 ml TID
*Cephalexin <i>Ulexin susp 60ml</i>	1.5g/60ml/bot	>25kg: 一般感染 : 250mg Q6H 嚴重感染 : 500~1000mg Q6H <25kg: 一般感染 : 25~50mg/kg/day Q6H 嚴重感染 : 50~100mg/kg/day Q6H
* Cetirizine HCl <i>Zyrtec</i>	1 mg/ml, 60 ml/bot	< 2 y/o: 0.25 ml/kg BID 2-6 y/o: 5 ml QD >6 y/o: 10 ml QD
Chloral hydrate 10% solution	2 mg/ml	<i>Sedation</i> : 25 mg/kg/day, Q8H <i>Hypnotic</i> : 50 mg/kg/dose
* Cyproheptadine HCl <i>Cyproh syrup</i>	0.4 mg /ml, 60 ml/bot	2-6 yrs: 5 ml TID 6-14yrs : 10 ml TID
* Digoxin <i>Cardiacin elixir</i>	50 mcg/ml, 60 ml/bot	10-20 mcg/kg/day
* Domperidone, <i>Wempty susp</i>	1 mg/ml, 60 ml/bot	0.25-0.5 ml/kg/dose TID-QID, before meal

台中榮總小兒科常用水劑藥品使用劑量表(II)

藥品名稱	成份含量	一般劑量
* <i>Ferrum Hausmann</i>	50 mg iron /ml (1drop = 2.5 mg iron), 30 ml/bot	<i>Infants:</i> Begin with 6 drops daily and gradually increase to 20 drops daily <i>Children:</i> 20 drops 1-2 times daily
<i>Fusoco syrup</i> 120 ml/bot	Each ml: Dextromethophan 0.5 mg Chlorpheniramine 0.15 mg <i>dl</i> -methylephedrine 0.4 mg Potassium guaiacol 6 mg Ammonium chloride 2 mg	2-4 yrs: 4 ml TID-QID 5-7 yrs: 7 ml /TID-QID 15-18 yrs : 10 ml /TID-QID
*Furosemide <i>Fumide oral solution</i>	10 mg/ml, 120 ml/bot	早產新生兒(<29 週): 1-2 mg/kg/dose Q24H 早產新生兒(>29 週): 1-2 mg/kg/dose Q12- 24H 新生兒: 1-3 mg/kg Q8H(需要時) 嬰兒及小孩: 起始劑量: 2 mg/kg/dose, 1-2 mg/kg/dose Q6-8H MAX 6 mg/kg/dose
Glyceryl Guaiacolate <i>Unitussin syrup</i>	20 mg/ml, 200 ml/bot	2-6 yrs : 2.5 ml QID 6-12 yrs : 5 ml QID
Ibuprofen <i>U-PEDIA</i>	20 mg/ml, 60 ml/bot	6 m/o-12 yrs: 0.25-0.5 ml/kg
Ketotifen <i>Athmin syrup</i>	0.2 mg/ml, 60ml/bot	0.5-2 mg BID
* Loratadin <i>Clarityne syrup</i>	5 mg/5 ml, 60 ml/bot	2-12 yrs:<30 kg: 5 mg QD >30 kg: 10 mg QD
* <i>Mycostatin susp</i>	100,000 U/ml, 24 ml/bot	1 ml QID (需要時可增加劑量)
<i>Neutral Phospate solution</i> powder/pack	Phosphorus: 200 mg/10ml Sodium: 1.08 mEq/ml 500 ml/bot	視需要適量投與
*Oxcarbazepine <i>Trileptal susp</i>	60 mg/ml, 100 ml/bot	4-16 yrs: 起始劑量: 8-10 mg/kg/day分2次 服用
<i>Pecolin susp</i> 120 ml/bot	Each 15ml: Kaolin 3 gm + Pectin 150 mg	3 ~ 6 yrs: 7 ml TID-QID >6 ~ 12 yrs:15 ml TID-QID
* Prednisolone <i>Kidsolone oral susp</i>	1 mg/ml, 60 ml/bot	遵照醫囑

台中榮總小兒科常用水劑藥品使用劑量表(III)

藥品名稱	成份含量	一般劑量
* Procatamol <i>Meptin solution</i>	5 mcg/ml, 60 ml/bot	< 6 y/o: 10-15 mcg QD > 6 y/o: 25 mcg QD-BID
* Simethicon <i>Wilcon susp</i>	20 mg/ml, 300 ml/bot	< 2 yrs : 1ml QID > 2 yrs : 2ml QID
* <i>SulfacoTrim syr</i> 60ml/bot	Each 5ml: Trimethoprin 40mg + Sulfamethoxazole 200mg	6-10 mg TMP/kg/day, Q12H
* Theophylline <i>Centertheo liquid</i>	5.34 mg/ml, 60 ml/bot	6-12 yrs: 125-250 mg Q6H (3 ml/kg/day Q8-6H or 1 ml/kg/dose)
* Valproate Sod. <i>Depakine solution</i>	200 mg/ml, 40 ml/bot	20-30 mg/kg/day, BID - TID

*有此星號標示之水劑為本院另有口服錠劑或膠囊劑型，因此對於較小的幼兒病患，請醫師們儘量開立水劑藥品

藥劑部 2011-03更新